

ABSTRACT

Objectives. The largest known dioxin contamination occurred between 1962 and 1970, when 12 million gallons of Agent Orange, a defoliant mixture contaminated with a form of the most toxic dioxin, were sprayed over southern and central Vietnam. Studies were performed to determine if elevated dioxin levels persist in Vietnamese living in the south of Vietnam.

Methods. With gas chromatography and mass spectroscopy, human milk, adipose tissue, and blood from Vietnamese living in sprayed and unsprayed areas were analyzed, some individually and some pooled, for dioxins and the closely related dibenzofurans.

Results. One hundred sixty dioxin analyses of tissue from 3243 persons were performed. Elevated 2,3,7,8-tetrachlorodibenzo-p-dioxin (TCDD) levels as high as 1832 ppt were found in milk lipid collected from southern Vietnam in 1970, and levels up to 103 ppt were found in adipose tissue in the 1980s. Pooled blood collected from southern Vietnam in 1991/92 also showed elevated TCDD up to 33 ppt, whereas tissue from northern Vietnam (where Agent Orange was not used) revealed TCDD levels at or below 2.9 ppt.

Conclusions. Although most Agent Orange studies have focused on American veterans, many Vietnamese had greater exposure. Because health consequences of dioxin contamination are more likely to be found in Vietnamese living in Vietnam than in any other populations, Vietnam provides a unique setting for dioxin studies. (*Am J Public Health*. 1995;85:516-522)

Agent Orange and the Vietnamese: The Persistence of Elevated Dioxin Levels in Human Tissues

Arnold Schecter, MD, MPH, Le Cao Dai, MD, Le Thi Bich Thuy, MPH, Hoang Trong Quynh, MD, Dinh Quang Minh, MD, Hoang Dinh Cau, MD, Pham Hoang Phiet, MD, Nguyen Thi Ngoc Phuong, MD, John D. Constable, MD, Robert Baughman, PhD, Olaf Pöpke, MS, J. J. Ryan, PhD, Peter Fürst, PhD, and Seppo Räsänen, PhD

Introduction

Studies of Agent Orange and its dioxin contaminant, 2,3,7,8-tetrachlorodibenzo-p-dioxin (TCDD), commonly referred to as "dioxin," have focused mainly on American veterans.¹ The health consequences for the Vietnamese have seldom been considered. This paper summarizes the assessments of exposure conducted from 1970 to 1974 and from 1983 to 1994.

Of the herbicides sprayed during the Vietnam War (known in Vietnam as the Second Indochina War), Agent Orange was by far the most abundant.^{1,2} Throughout the course of that war, which ended in 1975, American and South Vietnamese fixed-wing aircraft sprayed more than 12 million gallons of Agent Orange over about 10% of what was then South Vietnam, the area below the 17th parallel. Additional spraying was conducted from helicopters, from back-packs, and from boats. Spraying began in 1962, intensified in 1967, and is believed to have ended in 1971. This phenoxyherbicide defoliant, named for the orange coding stripes on the 55-gal barrels in which it was stored, consisted of 50% 2,4-dichlorophenoxyacetic acid (2,4-D) and 50% 2,4,5-trichlorophenoxyacetic acid (2,4,5-T) in an n-butyl ester formulation. The latter was contaminated with the most toxic dioxin congener, 2,3,7,8-TCDD, which is believed to have been present in Agent Orange at an average level of 3 ppm. In Agent Purple, a related herbicide, TCDD contamination averaged 30 ppm.

Because TCDD is very persistent in human tissue and the environment,¹⁻⁴ the potential health effects of Agent Orange are of particular concern to the Vietnamese people, some of whom have been at risk

since the spraying began in 1962. Eighty percent of the population live in rural areas. They traditionally wear open sandals or walk barefoot while working in the fields; they eat food grown on contaminated soil; and they consume water from contaminated areas. By contrast, American veterans generally served in Vietnam for only 1 year and consumed US-supplied food.


Exposure to Agent Orange can best be assessed by identifying tissue levels of 2,3,7,8-TCDD, the dioxin characteristic of

Arnold Schecter is with the Department of Preventive Medicine, Clinical Campus, State University of New York Health Science Center-Syracuse, Binghamton, NY. Le Cao Dai, Le Thi Bich Thuy, Hoang Trong Quynh, Dinh Quang Minh, and Hoang Dinh Cau are with the National Committee for Investigation of the Consequences of Chemicals Used in the Vietnam War, 10/80 Committee, at the College of Medicine, Hanoi, Viet Nam. Pham Hoang Phiet is with Cho Ray Hospital and Ho Chi Minh City Medical School, and Nguyen Thi Ngoc Phuong is with Tu Du Obstetrics and Gynecology Hospital and the College of Medicine, Ho Chi Minh City, Viet Nam. John D. Constable and Robert Baughman are with Harvard Medical School, Boston, Mass. Olaf Pöpke is with ERGO Forschungsgesellschaft mbH, Hamburg, Germany. J. J. Ryan is with the Food Research Division of Health Canada, Ottawa. Peter Fürst is with the State Laboratory for Food, Environmental and Pharmaceutical Chemistry, Münster, Germany. Seppo Räsänen is with the Chemical Instrument Center, University of Helsinki, Finland.

Requests for reprints should be sent to Arnold Schecter, MD, MPH, State University of New York Health Science Center-Syracuse, Clinical Campus, PO Box 1000, Binghamton, NY 13902.

This paper was accepted December 13, 1994.

Editor's Note. See related editorial by Dwyer and Flesch-Janys (p 476) in this issue.



1.9, and 0.6 ppt, respectively). In southern specimens from the 1980s, the highest individual TCDD level was 103 ppt. In a pooled adipose tissue sample from 10 northern soldiers who had served for many years in Agent Orange-sprayed jungles in the south, the TCDD level was 8.1 ppt (not shown on Table 1). In the northern samples from the general population, the highest level was 2.9 ppt.

Figure 1 illustrates TCDD levels in human milk over time. During the period in which fixed-wing aircraft sprayed Agent Orange and just after it ended in 1970, these levels were quite high; they then gradually declined. The highest dioxin level in 1970 specimens was approximately 1832 ppt, lipid.²⁵ Of individual samples from five women in 1970, positive levels shown varied from approximately 333 to 1832 ppt. There were also several samples in which TCDD was not detected. In three human milk samples obtained in 1973, TCDD levels ranged from 133 to 280 ppt, lipid. Several other samples had undetectable levels. Aliquots taken from 1973 archived milk previously analyzed by Baughman were found to contain TCDD levels between 77 and 230 ppt when analyzed by Ryan; these results were similar to the original data.

Table 2 shows dioxin contamination from Agent Orange as well as from other sources. In contrast to TCDD, found in Agent Orange, higher chlorinated dioxins and dibenzofurans with five to eight chlorines are found in chlorinated phenols used as wood preservatives and in agriculture. The mean results of pooled blood analyses by specific dioxin or dibenzofuran congener and by geographic region are presented for the first 698 patients in our pooled-blood series of 1991/92. These results are converted to dioxin toxic equivalents, reflecting total dioxin toxicity using "international" weighting toxic equivalency factors.³⁷⁻³⁹ As previously noted, these factors range from 1.0 for 2,3,7,8-TCDD to 0.001 for octachlorodibenzo dioxin. Here, mean TCDD level varies from 2.2 ppt in the north to 12.9 ppt and 13.2 ppt in the south and central regions, respectively. Total measured blood PCDDs and PCDFs, reflecting dioxins (and dibenzofurans) from industrial sources⁴⁰⁻⁴⁴ as well as Agent Orange, averaged 853 ppt in the south, 1145 ppt in the central region, and 287 ppt in the north. Total PCDD and PCDF dioxin toxic equivalents averaged 31.3, 50, and 15.3 in these blood samples.

Table 3 presents TCDD levels and total dioxin toxic equivalents in the 43

TABLE 2—Congener-Specific Dioxin Levels and Dioxin Toxic Equivalents (Lipid Basis, Parts per Trillion) for Pooled Vietnamese Blood, by Geographic Region, 1991 through 1992 Samples

| Congener | TEF | South (n = 433) | | Central (n = 183) | | North (n = 82) | |
|----------------------------------|-------|-----------------|-------|-------------------|-------|----------------|-------|
| | | Measured Level | TEq | Measured Level | TEq | Measured Level | TEq |
| Dioxins (PCDDs) | | | | | | | |
| 2,3,7,8-TCDD | 1.0 | 12.9 | 12.9 | 13.2 | 13.2 | 2.2 | 2.2 |
| 1,2,3,7,8-PeCDD | 0.5 | 8.0 | 4.0 | 16.3 | 8.2 | 4.1 | 2.1 |
| 1,2,3,4,7,8-HxCDD | 0.1 | 6.6 | 0.7 | 13.0 | 1.3 | 3.7 | 0.4 |
| 1,2,3,6,7,8-HxCDD | 0.1 | 29.9 | 2.9 | 46.2 | 4.6 | 13.4 | 1.3 |
| 1,2,3,7,8,9-HxCDD | 0.1 | 8.3 | 0.8 | 13.4 | 1.3 | 4.8 | 0.5 |
| 1,2,3,4,6,7,8-HpCDD | 0.01 | 77.2 | 0.8 | 78.1 | 0.8 | 25.5 | 0.3 |
| OCDD | 0.001 | 616.1 | 0.6 | 751.0 | 0.8 | 132.1 | 0.1 |
| Dibenzofurans (PCDFs) | | | | | | | |
| 2,3,7,8-TCDF | 0.1 | 2.1 | 0.2 | 2.9 | 0.3 | 4.6 | 0.5 |
| 1,2,3,7,8-PeCDF | 0.05 | 1.8 | 0.09 | 2.2 | 0.1 | 1.7 | 0.09 |
| 2,3,4,7,8-PeCDF | 0.5 | 8.3 | 4.2 | 14.9 | 7.5 | 7.6 | 3.8 |
| 1,2,3,4,7,8-HxCDF | 0.1 | 21.1 | 2.1 | 67.4 | 6.7 | 20.6 | 2.1 |
| 1,2,3,6,7,8-HxCDF | 0.1 | 13.0 | 1.3 | 40.0 | 4.0 | 11.1 | 1.1 |
| 1,2,3,7,8,9-HxCDF | 0.1 | 0.7 | 0.1 | 0.7 | 0.1 | 0.5 | 0.1 |
| 2,3,4,6,7,8-HxCDF | 0.1 | 2.3 | 0.2 | 3.0 | 0.3 | 2.2 | 0.2 |
| 1,2,3,4,6,7,8-HpCDF | 0.01 | 37.8 | 0.4 | 75.7 | 0.8 | 46.7 | 0.5 |
| 1,2,3,4,7,8,9-HpCDF | 0.01 | 3.4 | 0.03 | 1.9 | 0.02 | 1.9 | 0.02 |
| OCDF | 0.001 | 3.9 | 0.004 | 5.1 | 0.005 | 4.2 | 0.004 |
| Dioxins and dibenzofurans | | | | | | | |
| Total PCDDs | | 759.0 | 22.7 | 931.2 | 30.2 | 185.8 | 6.9 |
| Total PCDFs | | 94.4 | 8.6 | 213.8 | 19.8 | 101.1 | 8.4 |
| Total PCDD/Fs | | 853.4 | 31.3 | 1145.0 | 50 | 286.9 | 15.3 |

Note. TEF = dioxin toxic equivalency factor (toxicity relative to TCDD); TEq = dioxin toxic equivalent (measured dioxin level × TEF); n = number of pooled samples.

southern and central areas of Vietnam and 144 from the north).

In the southern samples, mean TCDD levels in blood, milk, and adipose

tissue are relatively similar (12.6, 7.5, and 14.7 ppt, lipid, respectively). In samples from the north, mean levels for blood, milk, and adipose tissue are lower (2.2,

These studies in Vietnam were conducted under less than ideal circumstances. During the early period, the country was involved in military conflicts, first with the United States (ending in 1975) and then with Cambodia and China. From 1978 to 1994, the country was under a US-imposed economic embargo. But even under these difficult conditions, the various human tissue samples, collected opportunistically and sporadically, clearly document elevated levels of 2,3,7,8-TCDD, the only dioxin contaminant of Agent Orange, at much higher levels in persons living in areas sprayed in southern Vietnam than in persons living in the unsprayed north, above the 17th parallel, where we find some of the lowest dioxin tissue levels reported worldwide to date.^{31,43}

In milk samples collected from nursing women in 1970, during the time of spraying, we found the highest dioxin levels reported in milk to date: approximately 1832 ppt TCDD.^{2,48} In samples collected in 1973, three years after the spraying of Agent Orange is thought to have ended, somewhat lower but also markedly elevated levels of dioxin were still found in milk. At present, levels in milk from the southern samples are declining to levels similar to those found in industrial countries, although they are still higher than those found in the north of Vietnam. In the United States, for example, TCDD levels are 3 to 6 ppt in the general population and total dioxin toxic equivalents are between 20 and 40 ppt, lipid.^{4,49}

Dioxins originate from many sources. These include municipal waste or toxic waste incineration, paper and pulp bleaching using chlorine, chlorinated phenols used as fungicides, wood preservatives and pesticides, feed stocks used in chemical production, herbicides, and polychlorinated biphenyl transformer fires.⁴⁰⁻⁴⁴ There is a characteristic pattern of dioxins and dibenzofurans from each of these sources, as there is for Agent Orange, where only TCDD is characteristic.

As noted previously, total dioxin toxicity, characterized by dioxin toxic equivalents from all congeners, is higher than toxicity of TCDD alone. This will become even more the case as dioxins, dibenzofurans, and dioxin-like polychlorinated biphenyls from industrial processes and agricultural use add to the human body burden of dioxin-like chemicals. Thus, accurate exposure assessments, whether for environmental fate, risk assessment, or dioxin health studies, need to

consider all dioxins present and total dioxin toxic equivalents, not just 2,3,7,8-TCDD. Blood values of TCDD may be elevated but total dioxin toxicity may not be, or the reverse might be the case. Both scenarios are seen in Table 3. Thus, total dioxin toxicity is not always proportionate to Agent Orange exposure.

Mapping the geographic areas where TCDD elevation in Vietnamese tissue reflects the presence and bioavailability of dioxin from Agent Orange can be helpful to others besides the Vietnamese. With areas of likely intake identified, large numbers of individual blood dioxin analyses of Vietnam veterans from the United States, Korea, and Australia might not be necessary.

Cost is an important consideration in planning research studies. Complete dioxin analyses of blood performed by one of the less than 30 World Health Organization certified laboratories currently cost up to \$2000 each. Collection, shipping, and medical interpretation further increase the cost. Studies using pooled blood data can rapidly and economically provide public health information on average population dioxin levels, despite certain methodological limitations.

Now that Vietnamese-US scientific teams are in place and have years of experience working together, environmental mapping of Agent Orange in Vietnam that uses 2,3,7,8-TCDD elevation relative to other dioxins in blood as a marker can be completed relatively quickly, given sufficient funding. This should pave the way for the important Agent Orange- and dioxin-related studies of health outcomes in Vietnam.

Understanding the health effects of dioxin exposure in Vietnam will be valuable, not only for the almost forgotten Vietnamese, but also for the United States and other industrialized nations that seek to evaluate the health risks of widespread exposure to TCDD and other PCDDs/PCDFs. The health of American veterans exposed to Agent Orange is of concern, as is the health of Vietnamese in the south, the population most at risk. In addition, there are about 1 million immigrants from Vietnam, Laos, and Cambodia now living in the United States; these persons might also benefit from research in Vietnam since they too are potentially at risk for adverse health effects of dioxins.

The US Environmental Protection Agency Draft Dioxin Reassessment Documents^{50,51} (released for public and scientific review in late 1994 and awaiting

finalization from the Science Advisory Board of EPA) conclude from an extensive review of dioxins' toxicity and of human exposure that levels of dioxins found in the general US population may be at or close to levels that have consequences for health. Possible consequences are increased risk for cancers, adverse reproductive and developmental effects, immune deficiency, endocrine disruption, neurological damage including cognitive and behavioral damage from in utero exposure, and other health effects.⁵² Since the dioxin levels shown here in Vietnam often exceed US levels, this suggests that health consequences are all the more likely to be expected in Agent Orange-exposed Vietnamese. □

Acknowledgments

This work was partially funded by a number of sources, which are gratefully acknowledged. They include the American Association for the Advancement of Science, the National Institute of Environmental Health Sciences, the Christopher Reynolds Foundation, the Samuel Rubin Foundation, the CS Fund and Warsh-Mott Legacy, the Kunstader Family Foundation, Church World Service, and CIDSE-Trócaire, The Catholic Agency for World Development.

Other contributing dioxin chemists for the Vietnam studies included B. Eitzer and R. Hites for sediment analyses, M. Gross and colleagues from the University of Nebraska at Lincoln for Vietnamese adipose tissue and sediment analyses, and K. Olie from the University of Amsterdam for environmental and food sample analyses. Many scientists and physicians in Vietnam cooperated in the collection of samples; they include Dr Duong Thi Cuong, director of the Obstetrics Hospital in Hanoi, Dr Huynh Kim Chi of Song Be Hospital in Song Be province, Professor Pham Ho, Dr Bui Thi Lang, and Madame Nguyen Phoc Dai. Participating hospitals include the Pediatrics Hospitals, Binh Dan Hospital, and Oncology Hospital of Ho Chi Minh City; Viet Duc Hospital, Oncology Hospital, and the Children's Hospitals in Hanoi; and hospitals in Hue and Da Nang. We also wish to thank the many Vietnamese volunteers and to express our concern for the many Vietnamese and Americans who suffered as a result of the Vietnam War, in which Agent Orange was but one hazard. Thanks are extended also to Martha Schechter for her editorial comments and suggestions, which were very helpful. The manuscript was prepared with the assistance of Ruth Stento and Heather Kessler.

References

1. Institute of Medicine. *Veterans and Agent Orange: Health Effects of Herbicides Used in Vietnam*. Washington, DC: National Academy Press; 1994.
2. Westing AH. Herbicides in war: past and present. In: Westing AH, ed. *Herbicides in War: The Long-term Ecological and Human Consequences*. London, England: Taylor & Francis; 1984:1-24.

- compounds in incinerator effluents: a review of measurements and mechanisms of formation. *Chemosphere*. 1980;9:501-522.
42. Clement RE, Suter SA, Reiner E, McCurvin D, Hollinger D. Concentrations of chlorinated dibenzo-p-dioxins and dibenzofurans in effluents and centrifuged particulates from Ontario pulp and paper mills. *Chemosphere*. 1989;19:649-654.
 43. Crosby DG, Moilanen KW, Wong AS. Environmental generation and degradation of dibenzodioxins and dibenzofurans. *Environ Health Perspect*. 1973;259-266.
 44. Hutzinger O, Blumich MJ, Berg MVD, Olie K. Sources and fates of PCDDs and PCDFs: an overview. *Chemosphere*. 1985;14: 581-600.
 45. Patterson DG, Fürst P, Henderson LO, et al. Partitioning of in vivo bound PCDDs/PCDFs among various compartments in whole blood. *Chemosphere*. 1989;19:135-142.
 46. Needham LL, Burse VW, Head SL, et al. Adipose tissue/serum partitioning of chlorinated hydrocarbon pesticides in humans. *Chemosphere*. 1990;20:975-980.
 47. Schechter AJ, Pöpke O, Ball M, Ryan JJ. Partitioning of dioxins and dibenzofurans: whole blood, blood plasma and adipose tissue. *Chemosphere*. 1991;23:1913-1919.
 48. Schechter AJ, Dekin A, Weerasinghe NCA, Arghestani S, Gross ML. Sources of dioxins in the environment: a study of PCDDs and PCDFs in ancient, frozen Eskimo tissue. *Chemosphere*. 1988;17:627-631.
 49. World Health Organization. *Levels of PCBs, PCDDs, and PCDFs in Breast Milk*. 34th ed. Copenhagen, Denmark: FADL Publishers; 1989. Environmental Health #34.
 50. *Health Assessment Document for 2,3,7,8-Tetrachlorodibenzo-p-dioxin (TCDD) and Related Compounds*. Washington, DC: US Environmental Protection Agency; released 1994. EPA/600/BP-92/001a. Draft Dioxin Reassessment Document.
 51. *Estimating Exposure to Dioxin-like Compounds*. Washington, DC: US Environmental Protection Agency; released 1994. EPA/600/6-88/005Cb. Draft Dioxin Reassessment Document.
 52. Schechter A, ed. *Dioxins and Health*. New York, NY: Plenum Press; 1994.